## Poster presentation

## **Open Access** Local planar dendritic structure: a uniquely biological phenomenon? Yihwa Kim<sup>\*1</sup>, Robert Sinclair<sup>2</sup> and Erik De Schutter<sup>1,3</sup>

Address: <sup>1</sup>Computational Neuroscience Unit, Okinawa Institute of Science and Technology, Okinawa 904-0411, Japan, <sup>2</sup>Mathematical Biology Unit, Okinawa Institute of Science and Technology, Okinawa 904-2234, Japan and <sup>3</sup>Theoretical Neurobiology, University of Antwerp, B-2610 Antwerpen, Belgium

Email: Yihwa Kim\* - ykim@oist.jp

\* Corresponding author

from Eighteenth Annual Computational Neuroscience Meeting: CNS\*2009 Berlin, Germany. 18-23 July 2009

Published: 13 July 2009

BMC Neuroscience 2009, 10(Suppl 1):P4 doi:10.1186/1471-2202-10-S1-P4

This abstract is available from: http://www.biomedcentral.com/1471-2202/10/S1/P4

© 2009 Kim et al; licensee BioMed Central Ltd.

Dendrites of different neurons have their own characteristic structures [1]. Global planar structures have been observed in Purkinje cells and retinal ganglion cells. Further, local planar dendritic structures have been discovered in the pyramidal neurons of the rabbit visual cortex [2]. Given an enclosing bundle of parallel fibers, the global planar dendritic structures of Purkinje cells can be explained as a minimization of total dendritic and path lengths (from synapses to soma) [3]. We investigated whether a similar minimization approach can explain local dendritic structures and conducted two types of Monte Carlo simulation studies of local dendritic angles. In the first study, the coordinates for each point were determined at random from a uniform distribution and then the resulting points were connected randomly. The mean of resulting cone angles was close to 90 degrees (Figure 1). In the second study, randomly chosen points were connected to minimize the local connection length. Interestingly, the resulting cone angle distribution had its peak close to 180 degrees (Figure 2), implying that the local planar structures observed in the biological system [2] can be obtained by minimization of local connection length. Furthermore, we discovered that essentially the same distribution of local dendritic angles as the one in Figure 2 could be achieved when a parent point located at the center of the sphere branches uniformly to the local area. This is due to the fact that there is a higher probability of sampling coordinates that result in wide angles. Hence, previously suggested mechanisms for local planar struc-

ture, such as tension or forced equilibria between the parent and daughter segments [5], should be revisited under the naturally occurring distribution (Figure 2). Besides, the finding that most neuron types indeed have local planar structures further supports our result.



## Figure I

Monte Carlo random sampling of 4 points in space. The cone angle between a parent and two children branches is depicted.



Figure 2

Monte Carlo simulation of 4 points with optimization on local connection lengths.

## References

- 1. Cajal SR: Histology of the nervous system of man and vertebrates Oxford Univ. Press, Oxford; 1995.
- Uylings HBM, Smit GJ: Three-dimensional branching structure of pyramidal cell dendrites. Brain Research 1974, 87:55-60.
- Wen Q, Chklovskii DB: A cost-benefit analysis of neuronal morphology. J Neurophysiol 2008, 99:2320-2328.
- 4. Cherniak C: Local optimization of neuron arbors. *Biol Cybern* 1992, **66:**503-510.
- 5. Uylings HBM, Van Pelt J: Measures for quantifying dendritic arborizations. Network: Comput Neural Syst 2002, 13:397-414.

